

Source: Okayama University (JAPAN), Public Relations and Information Strategy

For immediate release: 27 February 2017

Okayama University research: Potential cause of Lafora disease revealed

(Okayama, 27 February) **Researchers at Okayama University report in the journal *Matters*, a potential third factor in the pathogenesis of Lafora disease. In addition to gene mutations, chemical modification of the enzyme Laforin induced by nitrosative stress is found to be a likely cause of the illness too.**

Lafora disease is a fatal illness characterized by the abnormal generation of glycogens within neurons and other cells, for which no cure exists. Classified as a neurodegenerative disorder, it is caused by gene mutations that affect the production of two enzymes in the human body, Laforin and Malin. A team of researchers from Okayama University led by Ayano Satoh have now identified another potential cause of Lafora disease: when Laforin undergoes a particular reaction known as S-nitrosylation, the enzyme stops functioning properly.

The researchers reasoned that since S-nitrosylation has been linked to the development of other neurodegenerative diseases like Alzheimer's and Parkinson's, the reaction may also play a role in the pathogenesis of Laforin disease. They first confirmed that S-nitrosylation does take place by letting Laforin react with nitric oxide (NO). The authors then investigated whether the S-nitrosylation of Laforin affected its three biochemical functions.

One of Laforin's roles is to remove the phosphates that get incorporated, due to enzymatic errors, in glycogens. This is a three-step process: Laforin binds to the glycogen, removes the phosphate, and then binds to Malin. Satoh and colleagues checked whether S-nitrosylated Laforin still binds to glycogens and to Malin; their results suggest that modified Laforin has the same binding capacities as unmodified Laforin. Laforin's ability to remove phosphate groups — its phosphatase function — was, however, found to be inhibited after S-nitrosylation. The authors then looked at what happened when treating cells expressing non-mutated Laforin with an NO donor. A comparison with untreated cells revealed that 40% more Lafora bodies were formed when Laforin was treated with NO; Lafora bodies are the inclusion bodies characteristic of Lafora disease, consisting of accumulated glycogen with excessive phosphate content and precipitating inside cells.

The findings of Satoh and colleagues show that the decreased phosphatase activity of Laforin triggered by an overproduction of NO — as the result of a disrupted redox balance in cells — may be a cause, other than gene mutation, of Lafora disease. In the words of the researchers: "Lafora disease phenotypes can be caused by nitrosative stress". Additional research is needed for precisely determining Laforin's S-nitrosylation sites, and a further next step will be repeating the experiments in animal cells or neurons.

Background

Lafora disease


Lafora disease is a neurodegenerative disorder, like Alzheimer's, Parkinson's and Huntington's. In the case of Lafora disease, disordered cell metabolism leads to the accumulation of insoluble particles (called Lafora bodies) within the cytoplasm of cells in various parts of the human body. Symptoms appear in the patient's teenage years. Lafora disease is fatal; patients usually die within ten years after the first symptoms.

Known causes of the generation of Lafora bodies are mutations of genes that encode two enzymes, Laforin and Malin. The latter is a so-called E3 ubiquitin ligase: a protein that assists in transferring a ubiquitin molecule (a very common regulatory protein) from an E2 enzyme (an enzyme capable of carrying ubiquitin) to a substrate peptide. Laforin is a so-called glucan phosphatase: its function is to liberate phosphate groups from glucans (clusters consisting of a large number of glucose molecules). In the context of Lafora disease, the relevant glucans are glycogens (multi-branched strands of glucose molecules organized into a globular structure); Lafora bodies largely consist of phospho-glycogens. The presence of phosphates in glycogen makes it insoluble — hence the precipitation of Lafora bodies.

Satoh and colleagues hypothesized that not only mutations can cause Lafora disease, but also a certain type of chemical reaction involving Laforin: *S*-nitrosylation. This reaction modifies Laforin to such an extent that it can no longer perform its phosphate-liberating function. The researchers showed that Laforin molecules can indeed undergo *S*-nitrosylation reactions and that they lead to the increased formation of Lafora bodies in cells expressed with Laforin.

S-nitrosylation

By producing specific amounts of reactive oxygen species (ROS), reactive nitrogen species (RNS) and an antioxidant system, cells are able to maintain the redox-state balance necessary for proper metabolic functioning. In case of overproduction of ROS and RNS ('nitrosative stress'), reactions can be triggered that cause a disruption of cell metabolism. Nitric oxide (NO) is a ROS/RNS compound, and can react with Laforin; the reaction is known as *S*-nitrosylation. Because *S*-nitrosylation is recognized as a pathogenesis for Alzheimer's and Parkinson's diseases, the researchers proposed that it also plays a role in Laforin disease. They found that *S*-nitrosylated Laforin loses its phosphatase function, and that nitrosative stress may therefore be another cause of Lafora disease.



Caption

LB formation is increased by nitric oxide (NO) Representative images of (a) untreated or (b) NO donor treated wild-type Laforin-expressing cells are shown. Green, cells stained for Laforin and LB; blue, nuclei.

Reference

Rikako Toyota, Yasuko Honjo, Risa Imajo, Ayano Satoh. S-nitrosylation of laforin inhibits its phosphatase activity and is implicated in Lafora disease. *Matters*, Sep 8th, 2016.

DOI : 10.19185/matters.201606000014

<https://www.sciencematters.io/articles/201606000014>

Correspondence to

Associate Professor Ayano Satoh, Ph.D.
Organelle Systems Biotechnology,
Department of Medical Bioengineering, Graduate School of Natural
Science and Technology, Okayama University,
3-1-1 Tsushimanaka, Kita-ku, Okayama 700-8530, Japan
e-mail : ayano113@cc.okayama-u.ac.jp



Associate Professor Ayano Satoh

Further information

Okayama University
1-1-1 Tsushima-naka , Kita-ku , Okayama 700-8530, Japan
Public Relations and Information Strategy
E-mail: www-adm@adm.okayama-u.ac.jp
Website: http://www.okayama-u.ac.jp/index_e.html
Okayama Univ. e-Bulletin: <http://www.okayama-u.ac.jp/user/kouhou/ebulletin/>

Okayama Univ. e-Bulletin (PDF Issues): <http://www.okayama-u.ac.jp/en/tp/cooperation/ebulletin.html>

About Okayama University (You Tube):

<https://www.youtube.com/watch?v=iDL1coqPRYI>

Okayama University Image Movie (You Tube):

<https://www.youtube.com/watch?v=WnbJVk2eIA>

<https://www.youtube.com/watch?v=KU3hOIXS5kk>

Okayama University Medical Research Updates (OU-MRU)

Vol.1 : [Innovative non-invasive 'liquid biopsy' method to capture circulating tumor cells from blood samples for genetic testing](#)

Vol.2 : [Ensuring a cool recovery from cardiac arrest](#)

Vol.3 : [Organ regeneration research leaps forward](#)

Vol.4 : [Cardiac mechanosensitive integrator](#)

Vol.5 : [Cell injections get to the heart of congenital defects](#)

Vol.6 : [Fourth key molecule identified in bone development](#)

Vol.7 : [Anticancer virus solution provides an alternative to surgery](#)

Vol.8 : [Light-responsive dye stimulates sight in genetically blind patients](#)

Vol.9 : [Diabetes drug helps towards immunity against cancer](#)

Vol.10 : [Enzyme-inhibitors treat drug-resistant epilepsy](#)

Vol.11 : [Compound-protein combination shows promise for arthritis treatment](#)

Vol.12 : [Molecular features of the circadian clock system in fruit flies](#)

Vol.13 : [Peptide directs artificial tissue growth](#)

Vol.14 : [Simplified boron compound may treat brain tumours](#)

Vol.15 : [Metamaterial absorbers for infrared inspection technologies](#)

Vol.16 : [Epigenetics research traces how crickets restore lost limbs](#)

Vol.17 : [Cell research shows pathway for suppressing hepatitis B virus](#)

Vol.18 : [Therapeutic protein targets liver disease](#)

Vol.19 : [Study links signalling protein to osteoarthritis](#)

Vol.20 : [Lack of enzyme promotes fatty liver disease in thin patients](#)

Vol.21 : [Combined gene transduction and light therapy targets gastric cancer](#)

Vol.22 : [Medical supportive device for hemodialysis catheter puncture](#)

Vol.23 : [Development of low cost oral inactivated vaccines for dysentery](#)

Vol.24 : [Sticky molecules to tackle obesity and diabetes](#)

Vol.25 : [Self-administered aroma foot massage may reduce symptoms of anxiety](#)

Vol.26 : [Protein for preventing heart failure](#)

Vol.27 : [Keeping cells in shape to fight sepsis](#)

Vol.28 : [Viral-based therapy for bone cancer](#)

Vol.29 : [Photoreactive compound allows protein synthesis control with light](#)

Vol.30 : [Cancer stem cells' role in tumor growth revealed](#)

Vol.31 : [Prevention of RNA virus replication](#)

Vol.32 : [Enzyme target for slowing bladder cancer invasion](#)

Vol.33 : [Attacking tumors from the inside](#)

Vol.34 : [Novel mouse model for studying pancreatic cancer](#)

About Okayama University

Okayama University is one of the largest comprehensive universities in Japan with roots going back to the Medical Training Place sponsored by the Lord of Okayama and established in 1870. Now with 1,300 faculty and 14,000 students, the University offers courses in specialties ranging from medicine and pharmacy to humanities and physical sciences.

Okayama University is located in the heart of Japan approximately 3 hours west of Tokyo by Shinkansen.

